

Polymorphisms in genes related to one-carbon metabolism are not related to pancreatic cancer in PanScan and PanC4

Max Leenders · Samsiddhi Bhattacharjee · Paolo Vineis · Victoria Stevens · H. Bas Bueno-de-Mesquita · Xiao-Ou Shu · Laufey Amundadottir · Myron Gross · Geoffrey S. Tobias · Jean Wactawski-Wende · Alan A. Arslan · Eric J. Duell · Charles S. Fuchs · Steven Gallinger · Patricia Hartge · Robert N. Hoover · Elizabeth A. Holly · Eric J. Jacobs · Alison P. Klein · Charles Kooperberg · Andrea LaCroix · Donghui Li · Margaret T. Mandelson · Sara H. Olson · Gloria Petersen · Harvey A. Risch · Kai Yu · Brian M. Wolpin · Wei Zheng · Ilir Agalliu · Demetrius Albanes · Marie-Christine Boutron-Ruault · Paige M. Bracci · Julie E. Buring · Federico Canzian · Kenneth Chang · Stephen J. Chanock · Michelle Cotterchio · J. Michael Gaziano · Edward L. Giovannucci · Michael Goggins · Göran Hallmans · Susan E. Hankinson · Judith A. Hoffman-Bolton · David J. Hunter · Amy Hutchinson · Kevin B. Jacobs · Mazda Jenab · Kay-Tee Khaw · Peter Kraft · Vittorio Krogh · Robert C. Kurtz · Robert R. McWilliams · Julie B. Mendelsohn · Alpa V. Patel · Kari G. Rabe · Elio Riboli · Anne Tjønneland · Dimitrios Trichopoulos · Jarmo Virtamo · Kala Visvanathan · Joanne W. Elena · Herbert Yu · Anne Zeleniuch-Jacquotte · Rachael Z. Stolzenberg-Solomon

Received: 19 September 2012 / Accepted: 19 December 2012 / Published online: 19 January 2013
© Springer Science+Business Media Dordrecht 2013

Abstract

Purpose The evidence of a relation between folate intake and one-carbon metabolism (OCM) with pancreatic cancer (PanCa) is inconsistent. In this study, the association between genes and single-nucleotide polymorphisms (SNPs) related to OCM and PanCa was assessed.

Methods Using biochemical knowledge of the OCM pathway, we identified thirty-seven genes and 834 SNPs to examine in association with PanCa. Our study included 1,408 cases and 1,463 controls nested within twelve cohorts (PanScan). The ten SNPs and five genes with lowest *p* values (<0.02) were followed up in 2,323 cases and 2,340 controls from eight case–control studies (PanC4) that participated in PanScan2. The correlation of SNPs with metabolite levels was assessed for 649 controls from the

Electronic supplementary material The online version of this article (doi:10.1007/s10552-012-0138-0) contains supplementary material, which is available to authorized users.

M. Leenders · P. Vineis · E. Riboli
Department of Epidemiology and Biostatistics, School of Public Health, Imperial College, London, UK

M. Leenders (✉) · H. B. Bueno-de-Mesquita
Department of Gastroenterology and Hepatology, University Medical Center Utrecht, F02.618, Postal Box 85500, 3508GA Utrecht, The Netherlands
e-mail: M.Leenders-6@umcutrecht.nl

S. Bhattacharjee · L. Amundadottir · G. S. Tobias · P. Hartge · R. N. Hoover · K. Yu · D. Albanes · S. J. Chanock · A. Hutchinson · K. B. Jacobs · J. B. Mendelsohn · R. Z. Stolzenberg-Solomon
Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA

P. Vineis
HuGeF Foundation, Turin, Italy

V. Stevens · A. LaCroix · A. V. Patel
Department of Epidemiology, American Cancer Society, Atlanta, GA, USA

H. B. Bueno-de-Mesquita
National Institute for Public Health and the Environment, Bilthoven, The Netherlands

X.-O. Shu · W. Zheng
Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt-Ingram Cancer Center, Vanderbilt University, Nashville, TN, USA

L. Amundadottir · S. J. Chanock
Laboratory of Translational Genomics, Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA

European Prospective Investigation into Cancer and Nutrition.

Results When both stages were combined, we observed suggestive associations with PanCa for rs10887710 (*MATIA*) (OR 1.13, 95 %CI 1.04–1.23), rs1552462 (*SYT9*) (OR 1.27, 95 %CI 1.02–1.59), and rs7074891 (*CUBN*) (OR 1.91, 95 %CI 1.12–3.26). After correcting for multiple comparisons, no significant associations were observed in either the first or second stage. The three suggested SNPs showed no correlations with one-carbon biomarkers.

Conclusions This is the largest genetic study to date to examine the relation between germline variations in OCM-related genes polymorphisms and the risk of PanCa. Suggestive evidence for an association between polymorphisms and PanCa was observed among the cohort-nested studies, but this did not replicate in the case–control studies. Our results do not strongly support the hypothesis that genes related to OCM play a role in pancreatic carcinogenesis.

Keywords Pancreatic cancer · One-carbon metabolism · Polymorphisms · Biomarkers · Epidemiology

Introduction

Impaired DNA methylation is known to cause cancer through induction of chromosomal instability [1]. Folate and related nutrients (homocysteine, cysteine, methionine, cobalamin, and vitamin B6) are thought to influence

carcinogenesis through the one-carbon metabolism (OCM) pathway, which is involved in DNA repair, nucleotide synthesis, and methylation [2]. In particular, low folate levels may result in decreased *S*-adenosyl-methionine levels, the primary donor for methylation reactions. It has been hypothesized that hypomethylation may occur as result of altered folate metabolism [3]. One-carbon metabolites (i.e., folate and related nutrients) have been associated with several types of cancer [4–8], and inverse associations with pancreatic cancer (PanCa) have been observed for dietary folate, although study results have been inconsistent [9–16]. In addition, previous studies in association with PanCa for OCM-related SNPs observed a positive association with PanCa for the *MTHFR 677TT* genotype [17] and for two SNPs on the *MTRR* gene (rs162049 and rs10380) [18]. Genome-wide association studies (GWAS) of PanCa have not identified associations with genes involved with OCM [19–24].

In the present study, we used a two-stage approach to evaluate whether 31 genes thought to play a role in the OCM pathway (<http://wikipathways.org/index.php/Pathway:WP241>) were associated with PanCa. Additionally, genes (*ALPL*, *CPS1*, *CUBN*, *FUT2*, *PRICKLE2*, and *TCN1*) that included SNPs previously observed to predict one-carbon metabolite levels in GWAS [25–27] were also included. The first stage included nested case–control studies from twelve cohorts that participated in the National Cancer Institute (NCI) Pancreatic Cancer Cohort Consortium genome-wide association study (PanScan1) [19]. The second stage, used for replication, included eight case–control studies with

M. Gross
Department of Laboratory Medicine/Pathology, School of
Medicine, University of Minnesota, Minneapolis, MN, USA

J. Wactawski-Wende
Department of Social and Preventive Medicine, University at
Buffalo, State University of New York, Buffalo, NY, USA

A. A. Arslan
Department of Obstetrics and Gynecology, New York University
School of Medicine, New York, NY, USA

A. A. Arslan · A. Zeleniuch-Jacquotte
Department of Environmental Medicine, New York University
School of Medicine, New York, NY, USA

A. A. Arslan · A. Zeleniuch-Jacquotte
New York University Cancer Institute, New York, NY, USA

E. J. Duell
Catalan Institute of Oncology (ICO-IDIBELL), Barcelona, Spain

C. S. Fuchs · B. M. Wolpin
Department of Medical Oncology, Dana-Farber Cancer Institute,
Boston, MA, USA

C. S. Fuchs · B. M. Wolpin · E. L. Giovannucci ·
S. E. Hankinson · D. J. Hunter
Channing Laboratory, Department of Medicine, Brigham and
Women's Hospital and Harvard Medical School, Boston, MA,
USA

S. Gallinger
Samuel Lunenfeld Research Institute, Mount Sinai Hospital,
University of Toronto, Toronto, Canada

E. A. Holly · P. M. Bracci
Department of Epidemiology and Biostatistics, University
of California San Francisco, San Francisco, CA, USA

E. J. Jacobs
Epidemiology Research Program, American Cancer Society,
Atlanta, GA, USA

A. P. Klein
Department of Oncology, Johns Hopkins University School
of Medicine, Baltimore, MD, USA

GWAS data that participated in the Pancreatic Cancer Case–Control Consortium (PanC4) and were part of PanScan 2 [22].

Methods

The methods for PanScan1 and PanScan2 are described elsewhere [19, 22]. The study population included PanCa cases and control participants from the previously conducted GWAS in the PanScan1 and PanC4 [19, 22]. The first stage of our analyses included 1,408 incident cases and 1,463 controls from the twelve cohort studies that participated in PanScan1 [19]. The second stage included 2,323 cases and 2,340 controls from eight case–control studies that participated in PanC4 and were part of PanScan1 and PanScan2 [19, 22]. All included studies had available dietary and lifestyle data. Cases were defined as participants diagnosed with primary adenocarcinoma of the exocrine pancreas, and controls were matched to cases according to birth year, sex, and self-reported race/ethnicity and were free of PanCa at the time of recruitment [19, 22]. Genotyping was performed by the NCI Core Genotyping Facility using the Illumina HumanHap550 and HumanHap550-Duo SNP arrays (PanScan1) and Illumina Human 610-Quad arrays (PanScan2) [19, 22]. Each individual study obtained informed consent from study participants and approval from local institutional review board. The Special Studies Institutional Review Board of the NCI approved the pooled PanScan study [19, 22].

Details of the collection of data across the cohorts are described elsewhere [19]. A total of 37 genes and 834 SNPs were included in the study. From all studies, data on the participant's age and gender were obtained. In addition, five effective principal components or eigenvectors determined from the GWAS data were included as quantitative covariates to correct for population structure.

Data on one-carbon metabolite biomarkers (homocysteine, cysteine, methionine, folate, cobalamin, and vitamin B6) measured in plasma were available for 327 controls of the case–control study nested within the EPIC cohort. Biomarker levels were analyzed by MS-based methods in the laboratory of BEVITAL AS (<http://www.bevital.no>) in Bergen, Norway [28].

In the first stage, the association for each SNP with PanCa was examined within the cohorts of PanScan using unconditional logistic regression, adjusted for age and sex. To correct for population stratification, analyses were also adjusted for cohort and five principal components of population stratification. Allelic additive models were used where the risk allele was the least frequent allele. *p* values were adjusted for multiple testing using the permutation-based closed step-down minP procedure [29]. Genes were evaluated separately using the adaptive rank truncated product (ARTP) method [30] with 20 truncation points, using 10,000 permutations. This captures potential multiple association signals within a gene, accounting for gene size and LD structure. Gene ARTP *p* values were adjusted for multiple comparisons using a false discovery rate [31].

The ten SNPs and five genes that showed lowest *p* values for the association with PanCa in the cohort data were

A. P. Klein
Department of Epidemiology, Johns Hopkins Medical Institutions, The Bloomberg School of Public Health, The Sol Goldman Pancreatic Research Center, Baltimore, MD, USA

C. Kooperberg · M. T. Mandelson
Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA, USA

D. Li
Department of Gastrointestinal Medical Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX, USA

M. T. Mandelson
Group Health Center for Health Studies, Seattle, WA, USA

S. H. Olson
Department of Epidemiology and Biostatistics, Memorial Sloan-Kettering Cancer Center, New York, NY, USA

G. Petersen · K. G. Rabe
Department of Health Sciences Research, Mayo Clinic, Rochester, MN, USA

H. A. Risch · H. Yu
Yale University School of Public Health, New Haven, CT, USA

I. Agalliu
Department of Epidemiology and Population Health, Albert Einstein College of Medicine, Bronx, NY, USA

M.-C. Boutron-Ruault
INSERM, Institut Gustave-Roussy, Paris-Sud University, Villejuif, France

J. E. Buring
Divisions of Preventive Medicine and Aging, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA

J. E. Buring
Department of Ambulatory Care and Prevention, Harvard Medical School, Boston, MA, USA

F. Canzian
Division of Cancer Epidemiology, German Cancer Research Center (DFKZ), Heidelberg, Germany

followed up in eight case–control studies from PanC4, using similar analyses. Because fewer comparisons were performed in this second stage, a Bonferroni correction was applied to both SNP and gene p values to adjust for multiple comparisons. Finally, a combined analysis of both stages was performed. Because combined results were mainly driven by cohort results, these were adjusted for multiple comparisons using similar techniques as applied to the first stage.

Adaptive rank truncated product p values also were computed for the total OCM pathway in all stages using the top 5 genes as truncation points. An adjusted p value below 0.05 was considered statistically significant.

Modification of the top ten SNPs by dietary intake of folate (in tertiles) was assessed in the cohorts by testing the statistical significance of the multiplicative interaction terms using likelihood ratio tests. Differences in least-squares means for plasma metabolite levels (homocysteine, cysteine, methionine, folate, cobalamin, and vitamin B6) were compared per genotype for each SNP by a pairwise t test in controls only.

Results

In the first stage of our analyses (cohort studies), rs10887710 (*MATIA*) showed the lowest p value of all tested SNPs (allelic OR 1.24, 95 % CI 1.08–1.41, p 0.002)

(Table 1). However, after correcting for multiple comparisons, none of the SNPs were statistically significant. None of the top ten SNPs for the first stage showed a relation with PanCa in the second stage (case–control studies), with all p values greater than 0.2. When all studies were combined (cohort and case–control), statistically significant associations were found for rs10887710 (*MATIA*), rs1552462 (*SYT9*), and rs7074891 (*CUBN*). However, after correcting for multiple comparisons, the associations were not statistically significant.

Two genes (*MATIA*, p 0.021 and *TYMS*, p 0.033) showed statistically significant ARTP p values in the first stage (Table 2). In the replication stage, none of the genes showed an association with PanCa. When both stages were combined, *MTRR* (p for cohort stage: 0.19) showed a significant p value (p 0.05). After adjustment for multiple comparisons, no significant associations were found in all stages. Additionally, the pathway analysis showed ARTP-adjusted p values above 0.50 in all stages.

There were no significant interactions of the top one-carbon pathway SNPs or genes by dietary folate. Within the subset of EPIC participants with the one-carbon biomarkers, several associations were observed between one-carbon biomarkers and the ten SNPs with the lowest p values in the cohort stage. For three SNPs in *TYMS* (rs3819101, rs11873007, and rs3786355) that were in high linkage disequilibrium, the TT genotype correlated with higher

K. Chang
Comprehensive Digestive Disease Center, University of California, Irvine Medical Center, Orange, CA, USA

M. Cotterchio
Dalla Lana School of Public Health, University of Toronto, Toronto, ON, Canada

M. Cotterchio
Prevention and Cancer Control, Cancer Care Ontario, Toronto, ON, Canada

J. M. Gaziano
Physicians' Health Study, Divisions of Aging, Cardiovascular Medicine, and Preventive Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA

J. M. Gaziano
Veterans Affairs Boston Healthcare System, Massachusetts Veterans Epidemiology Research and Information Center, Boston, MA, USA

E. L. Giovannucci · S. E. Hankinson · D. J. Hunter · P. Kraft · D. Trichopoulos
Department of Epidemiology, Harvard School of Public Health, Boston, MA, USA

E. L. Giovannucci
Department of Nutrition, Harvard School of Public Health, Boston, MA, USA

M. Goggins
Departments of Oncology, Pathology and Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, USA

G. Hallmans
Department of Public Health and Clinical Medicine, Nutritional Research, Umeå University, Umeå, Sweden

J. A. Hoffman-Bolton · K. Visvanathan
Department of Epidemiology, The Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD, USA

A. Hutchinson · K. B. Jacobs
Core Genotyping Facility, Advanced Technology Program, SAIC-Frederick Inc., NCI-Frederick, Frederick, MD, USA

K. B. Jacobs
Bioinformed Consulting Services, Gaithersburg, MD, USA

M. Jenab
International Agency for Research on Cancer (IARC/WHO), Lyon, France

methionine levels than the CT (28.46 vs. 25.34 $\mu\text{mol/L}$; p 0.01) or CC genotype (25.45 $\mu\text{mol/L}$; p 0.01). The AA genotype at rs1835898 (*PRICKLE2*) was associated with higher folate (17.26 vs. 14.55 nmol/L for AG; p 0.04) and higher pyridoxal phosphate (PLP) concentrations (63.44 vs. 44.45 nmol/L for AG; p 0.02 and 41.35 nmol/L for GG, p 0.02), when compared to the AG or GG genotypes. The AC genotype at rs222338 (*PRICKLE2*) was associated with higher cobalamin levels than the CC genotype (477.45 vs. 379.26 pmol/L; p 0.03) (data not shown).

Discussion

This is the largest genetic study to date examining the relation between germline variations in OCM-related genes and PanCa risk. In the first stage of this study (cohort studies), evidence for an association between genes (*MAT1A* and *TYMS*) and SNPs (most notably rs10887710 [*MAT1A*]) involved in OCM with PanCa was observed. However, after correcting for multiple comparisons, these associations were not statistically significant, nor did the suggestive evidence replicate in the second stage (case–control studies) of this study. No interaction between dietary folate and any of the ten SNPs with lowest p values in the first stage was observed.

For several of the top ten SNPs, a correlation with one-carbon biomarkers was observed. This indicates that the SNPs possibly directly influence biomarker levels, which could strengthen evidence for a relation between these SNPs and PanCa. However, none of the SNPs replicated in a second stage and correlations with biomarkers could only be calculated from a subset of all included participants. Additionally, no clear association between one-carbon metabolites and risk of PanCa has been observed within EPIC using the same data on metabolites and PanCa [32].

The reason for observing no replication of the suggestive evidence observed in the first stage of the study may have been due to differences in study designs, as cohort studies may be less prone to survival bias. However, considering the large number of SNPs tested, it is most likely that the significant SNPs observed in the cohort studies, which did not replicate in case–control studies, were false-positives in the first stage of the analysis.

In conclusion, we observed suggestive associations between SNPs and genes involved in one-carbon metabolism and risk of PanCa in the first stage of this study (cohort studies), but none of the results replicated in the second stage (case–control studies). Our results do not strongly support the hypothesis that genes related to OCM play a role in pancreatic carcinogenesis.

K.-T. Khaw
Department of Public Health and Primary Care, Clinical Gerontology, Addenbrooke's Hospital, University of Cambridge, Cambridge, UK

P. Kraft
Department of Biostatistics, Harvard School of Public Health, Boston, MA, USA

V. Krogh
Nutritional Epidemiology Unit, Fondazione Istituto di Recovero e Cura a Carattere Scientifico (IRCCS) Istituto Nazionale dei Tumori, Milan, Italy

R. C. Kurtz
Department of Medicine, Memorial Sloan-Kettering Cancer Center, New York, NY, USA

R. R. McWilliams
Division of Medical Oncology, Mayo Clinic, Rochester, MN, USA

A. Tjønneland
Institute of Cancer Epidemiology, Danish Cancer Society, Copenhagen, Denmark

D. Trichopoulos
Bureau of Epidemiologic Research, Academy of Athens, Athens, Greece

J. Virtamo
Department of Chronic Disease Prevention, National Institute for Health and Welfare, Helsinki, Finland

J. W. Elena
Division of Cancer Prevention and Population Control, Department of Health and Human Services, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA

Table 1 Top ten SNPs from cohort with corresponding odds ratios and *p* values for cohort, case–control, and combined data

Marker ^a , Alleles ^b , Chr ^c , Location ^c , and Gene ^d	Subset ^e	MAF ^f	Subjects ^g	<i>p</i> value ^h	<i>p</i> value (adjusted for multiple testing) ⁱ	OR (95 % CI) ^j
rs10887710 (T,C), 10q23.1 (82019765), <i>MAT1A</i>	Cohort	0.17 0.20	1,463 1,407	0.002	0.564	1.24 (1.08–1.41)
	Replication, case–control	0.19 0.20	2,340 2,323	0.341	1.000	1.06 (0.94–1.18)
	Combined	0.18 0.20	3,803 3,730	0.005	0.990	1.13 (1.04–1.23)
rs1552462 (C,T), 11p15.4 (7213717), <i>SYT9</i>	Cohort	0.02 0.03	1,463 1,408	0.010	1.000	1.58 (1.12–2.23)
	Replication, case–control	0.02 0.03	2,338 2,320	0.600	1.000	1.08 (0.81–1.44)
	Combined	0.02 0.03	3,801 3,728	0.033	1.000	1.27 (1.02–1.59)
rs3819101 (C,T), 18p11.32 (667240), <i>TYMS</i>	Cohort	0.29 0.33	1,462 1,407	0.011	1.000	1.16 (1.03–1.30)
	Replication, case–control	0.32 0.33	2,336 2,323	0.784	1.000	1.01 (0.92–1.11)
	Combined	0.31 0.33	3,798 3,729	0.098	1.000	1.06 (0.99–1.14)
rs2276433 (C,T), 11p15.1 (20630443), <i>SLC6A5</i>	Cohort	0.38 0.41	1,463 1,407	0.013	1.000	1.15 (1.03–1.28)
	Replication, case–control	0.42 0.39	2,340 2,323	0.986	1.000	1.00 (0.89–1.12)
	Combined	0.40 0.40	3,803 3,730	0.800	1.000	0.99 (0.93–1.06)
rs11873007 (C,T), 18p11.32 (670380), <i>TYMS</i>	Cohort	0.30 0.33	1,359 1,302	0.013	1.000	1.16 (1.03–1.30)
	Replication, case–control	0.32 0.34	2,326 2,309	0.742	1.000	1.02 (0.93–1.12)
	Combined	0.31 0.34	3,685 3,611	0.100	1.000	1.06 (0.99–1.14)
rs7074891 (T,C), 10p13 (17228480), <i>CUBN</i>	Cohort	0.00 0.01	1,463 1,408	0.014	1.000	2.36 (1.22–4.55)
	Replication, case–control	0.01 0.01	2,340 2,322	0.356	1.000	1.45 (0.59–3.58)
	Combined	0.01 0.01	3,803 3,730	0.017	1.000	1.91 (1.12–3.26)
rs1835898 (A,G), 3p14.1 (64096330), <i>PRICKLE2</i>	Cohort	0.45 0.42	1,462 1,408	0.015	1.000	0.88 (0.79–0.98)
	Replication, case–control	0.44 0.44	2,339 2,323	0.748	1.000	1.01 (0.93–1.11)
	Combined	0.44 0.43	3,801 3,731	0.172	1.000	0.95 (0.89–1.02)
rs3740873 (G,T), 11p15.1 (20626016), <i>SLC6A5</i>	Cohort	0.45 0.49	1,462 1,407	0.016	1.000	1.14 (1.02–1.27)
	Replication, case–control	0.49 0.47	2,340 2,323	0.201	1.000	0.95 (0.87–1.03)
	Combined	0.47 0.48	3,802 3,730	0.646	1.000	1.02 (0.95–1.09)
rs3786355 (C,T), 18p11.32 (671962), <i>TYMS</i>	Cohort	0.30 0.33	1,463 1,408	0.016	1.000	1.15 (1.03–1.29)
	Replication, case–control	0.32 0.33	2,338 2,320	0.822	1.000	1.01 (0.92–1.11)
	Combined	0.31 0.33	3,801 3,728	0.125	1.000	1.06 (0.98–1.14)
rs222338 (C,A), 3p14.1 (64202692), <i>PRICKLE2</i>	Cohort	0.05 0.04	1,463 1,408	0.017	1.000	0.73 (0.57–0.95)
	Replication, case–control	0.05 0.06	2,334 2,320	0.751	1.000	1.03 (0.85–1.26)
	Combined	0.05 0.05	3,797 3,728	0.266	1.000	0.92 (0.79–1.07)

The SNP-level results from the unconditional logistic regression of the genotypes generated in a total of 1,407 pancreatic cancer cases and 1,463 controls

The analysis adjusted for age in ten-year categories, sex, study, five principal components of population stratification

OR Allelic odds ratio, CI 95 % confidence interval

^a NCBI dbSNP identifier

^b Major allele, minor allele

^c Chromosome and NCBI Human genome Build 36 location

^d Gene neighborhood within 20 kb upstream and 10 kb downstream of SNP

^e Subset: cohort, cohort studies; case–control, case–control studies; combined, all studies

^f Minor allele frequency in control and case participants

^g Controls, cases

^h 1 d.f. Wald test

ⁱ *P* values of cohort and combined stage adjusted for multiple testing by the step-down minP procedure. *p* values of case–control stage adjusted by Bonferroni

^j Estimate assuming additive model

Table 2 Top five genes from the cohort, with corresponding minP *p* values for cohort, case–control, and combined data, and other genes included in the pathway analysis

Gene ^a and Chr ^b	Subset ^c	<i>p</i> value (ARTP)	<i>p</i> value (adjusted)
<i>MAT1A</i> , 10q23.1	Cohort	0.021	0.607
	Replication, case–control	0.255	1.000
	Combined	0.275	0.849
<i>TYMS</i> , 18p11.32	Cohort	0.033	0.607
	Replication, case–control	0.377	1.000
	Combined	0.815	1.000
<i>GNMT</i> , 6p21.1	Cohort	0.132	0.863
	Replication, case–control	0.863	1.000
	Combined	0.832	1.000
<i>FUT2</i> , 19q13.33	Cohort	0.150	0.863
	Replication, case–control	0.546	1.000
	Combined	0.431	1.000
<i>MTRR</i> , 5p15.31	Cohort	0.188	0.863
	Replication, case–control	0.094	0.471
	Combined	0.048	0.595
<i>SLC6A5</i> , 11p15.1	Cohort	0.194	0.863
<i>MTHFS</i> , 15q25.1	Cohort	0.225	0.863
<i>MTHFR</i> , 1p36.3	Cohort	0.230	0.863
<i>FOLH1</i> , 11p11.2	Cohort	0.322	0.863
<i>SLC6A9</i> , 1p33	Cohort	0.333	0.863
<i>GCSH</i> , 16q23.2	Cohort	0.342	0.863
<i>TCN2</i> , 22q12.2	Cohort	0.357	0.863
<i>CBS</i> , 21q22.3	Cohort	0.404	0.863
<i>PRICKLE2</i> , 3p14.1	Cohort	0.405	0.863
<i>NBPF3</i> , 1p36.12	Cohort	0.407	0.863
<i>AMD1</i> , 6q21	Cohort	0.430	0.863
<i>CTH</i> , 1p31.1	Cohort	0.432	0.863
<i>MAT2A</i> , 2p11.2	Cohort	0.441	0.863
<i>GIF</i> , 11q13	Cohort	0.500	0.863
<i>CPS1</i> , 2q35	Cohort	0.504	0.863
<i>SHMT1</i> , 17p11.2	Cohort	0.536	0.863
<i>FOLR2</i> , 11q13.3–q13.5	Cohort	0.553	0.863
<i>MTHFD1</i> , 14q24	Cohort	0.564	0.863
<i>SLC1A5</i> , 19q13.3	Cohort	0.590	0.863
<i>GGH</i> , 8q12.3	Cohort	0.623	0.863
<i>ALDH1L1</i> , 3q21.3	Cohort	0.644	0.863
<i>BHMT</i> , 5q13.1–q15	Cohort	0.665	0.863
<i>FOLR3</i> , 11q13	Cohort	0.675	0.863
<i>SHMT2</i> , 12q12–q14	Cohort	0.683	0.863
<i>SYT9</i> , 11p15.4	Cohort	0.700	0.863
<i>CUBN</i> , 10p12.31	Cohort	0.724	0.865
<i>ALPL</i> , 1p36.12	Cohort	0.766	0.886

Table 2 continued

Gene ^a and Chr ^b	Subset ^c	<i>p</i> value (ARTP)	<i>p</i> value (adjusted)
<i>TCN1</i> , 11q11–q12	Cohort	0.826	0.901
<i>SLC19A1</i> , 21q22.3	Cohort	0.828	0.901
<i>MTR</i> , 1q43	Cohort	0.879	0.929
<i>DHFR</i> , 5q11.2–q13.2	Cohort	0.925	0.951
<i>GSS</i> , 20q11.2	Cohort	0.983	0.983

The gene-level results using the Adaptive Rank Truncated Product approach, of the 37 genes in a total of 1,407 pancreatic cancer cases and 1,463 controls. For each SNP, the logistic regression analysis used adjustment for age in 10-year categories, sex, study, five principal components of population stratification. Adjustment for multiple comparisons was done using an FDR correction for the cohort and combined stage, and using a Bonferroni correction for the case–control stage

^a Gene neighborhood within 20 kb upstream and 10 kb downstream of SNP

^b Chromosome and NCBI Human genome Build 36 location

^c Subset: cohort, cohort studies; replication, case–control, case–control studies; combined, all studies

Acknowledgments This research was supported by the Intramural Research Program of the National Institutes of Health, Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Department of Health and Human Services. This study has been funded by the World Cancer Research Fund (Grant No. 2008/51 to PV). For full acknowledgments, please see electronic supplementary material.

Conflict of interest None.

References

- Ehrlich M (2002) DNA methylation in cancer: too much, but also too little. *Oncogene* 21(35):5400–5413
- Tibbetts AS, Appling DR (2010) Compartmentalization of Mammalian folate-mediated one-carbon metabolism. *Annu Rev Nutr* 30:57–81
- Liu JJ, Ward RL (2010) Folate and one-carbon metabolism and its impact on aberrant DNA methylation in cancer. *Adv Genet* 71:79–121
- Larsson SC, Orsini N, Wolk A (2010) Vitamin B6 and risk of colorectal cancer: a meta-analysis of prospective studies. *JAMA* 303(11):1077–1083
- Kim D-H, Smith-Warner SA, Spiegelman D et al (2010) Pooled analyses of 13 prospective cohort studies on folate intake and colon cancer. *Cancer Causes Control* 21(11):1919–1930
- Collin SM, Metcalfe C, Refsum H et al (2010) Circulating folate, vitamin B12, homocysteine, vitamin B12 transport proteins, and risk of prostate cancer: a case-control study, systematic review, and meta-analysis. *Cancer Epidemiol Biomarkers Prev* 19(6):1632–1642
- Larsson SC, Giovannucci E, Wolk A (2007) Folate and risk of breast cancer: a meta-analysis. *J Natl Cancer Inst* 99(1):64–76
- Kim YI (1999) Folate and carcinogenesis: evidence, mechanisms, and implications. *J Nutr Biochem* 10(2):66–88

9. Baghurst PA, McMichael AJ, Slavotinek AH, Baghurst KI, Boyle P, Walker AM (1991) A case-control study of diet and cancer of the pancreas. *Am J Epidemiol* 134(2):167–179
10. Bravi F, Polesel J, Bosetti C et al (2011) Dietary intake of selected micronutrients and the risk of pancreatic cancer: an Italian case-control study. *Ann Oncol* 22(1):202–206
11. Gong Z, Holly EA, Bracci PM (2009) Intake of folate, vitamins B6, B12 and methionine and risk of pancreatic cancer in a large population-based case-control study. *Cancer Causes Control* 20(8):1317–1325
12. Larsson SC, Håkansson N, Giovannucci E, Wolk A (2006) Folate intake and pancreatic cancer incidence: a prospective study of Swedish women and men. *J Natl Cancer Inst* 98(6):407–413
13. Oaks BM, Dodd KW, Meinhold CL, Jiao L, Church TR, Stolzenberg-Solomon RZ (2010) Folate intake, post-folic acid grain fortification, and pancreatic cancer risk in the Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial. *Am J Clin Nutr* 91(2):449–455
14. Stolzenberg-Solomon RZ, Pietinen P, Barrett MJ, Taylor PR, Virtamo J, Albanes D (2001) Dietary and other methyl-group availability factors and pancreatic cancer risk in a cohort of male smokers. *Am J Epidemiol* 153(7):680–687
15. Silverman DT, Swanson CA, Gridley G et al (1998) Dietary and nutritional factors and pancreatic cancer: a case-control study based on direct interviews. *J Natl Cancer Inst* 90(22):1710–1719
16. Keszei AP, Verhage BAJ, Heinen MM, Goldbohm RA, van den Brandt PA (2009) Dietary folate and folate vitamins and the risk of pancreatic cancer in the Netherlands cohort study. *Cancer Epidemiol Biomarkers Prev* 18(6):1785–1791
17. Larsson SC, Giovannucci E, Wolk A (2006) Folate intake, MTHFR polymorphisms, and risk of esophageal, gastric, and pancreatic cancer: a meta-analysis. *Gastroenterology* 131(4):1271–1283
18. Ohnami S, Sato Y, Yoshimura K et al (2008) His595Tyr polymorphism in the methionine synthase reductase (MTRR) gene is associated with pancreatic cancer risk. *Gastroenterology* 135(2):477–488
19. Amundadottir L, Kraft P, Stolzenberg-Solomon RZ et al (2009) Genome-wide association study identifies variants in the ABO locus associated with susceptibility to pancreatic cancer. *Nat Genet* 41(9):986–990
20. Diergaarde B, Brand R, Lamb J et al (2010) Pooling-based genome-wide association study implicates gamma-glutamyl-transferase 1 (GGT1) gene in pancreatic carcinogenesis. *Pancreatology* 10(2–3):194–200
21. Low S-K, Kuchiba A, Zembutsu H et al (2010) Genome-wide association study of pancreatic cancer in Japanese population. *PLoS ONE* 5(7):e11824
22. Petersen GM, Amundadottir L, Fuchs CS et al (2010) A genome-wide association study identifies pancreatic cancer susceptibility loci on chromosomes 13q22.1, 1q32.1 and 5p15.33. *Nat Genet* 42(3):224–228
23. Willis JA, Olson SH, Orlov I et al (2012) A replication study and genome-wide scan of single-nucleotide polymorphisms associated with pancreatic cancer risk and overall survival. *Clin Cancer Res* 18(14):3942–3951
24. Wu C, Miao X, Huang L et al (2012) Genome-wide association study identifies five loci associated with susceptibility to pancreatic cancer in Chinese populations. *Nat Genet* 44(1):62–66
25. Tanaka T, Scheet P, Giusti B et al (2009) Genome-wide association study of vitamin B6, vitamin B12, folate, and homocysteine blood concentrations. *Am J Hum Genet* 84(4):477–482
26. Lange LA, Croteau-Chonka DC, Marvelle AF et al (2010) Genome-wide association study of homocysteine levels in Filipinos provides evidence for CPS1 in women and a stronger MTHFR effect in young adults. *Hum Mol Genet* 19(10):2050–2058
27. Hazra A, Kraft P, Lazarus R et al (2009) Genome-wide significant predictors of metabolites in the one-carbon metabolism pathway. *Hum Mol Genet* 18(23):4677–4687
28. Midttun Ø, Hustad S, Ueland PM (2009) Quantitative profiling of biomarkers related to B-vitamin status, tryptophan metabolism and inflammation in human plasma by liquid chromatography/tandem mass spectrometry. *Rapid Commun Mass Spectrom* 23(9):1371–1379
29. Westfall PH, Young SS (1993) Resampling-based multiple testing: examples and methods for p-value adjustment. Wiley, New York, NY
30. Yu K, Li Q, Bergen AW et al (2009) Pathway analysis by adaptive combination of P-values. *Genet Epidemiol* 33(8):700–709
31. Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol* 57(1):289–300
32. Chuang S-C, Stolzenberg-Solomon R, Ueland PM et al (2011) A U-shaped relationship between plasma folate and pancreatic cancer risk in the European Prospective Investigation into Cancer and Nutrition. *Eur J Cancer* 47(12):1808–1816